

A note on numerical solution of the parabolic-Schrödinger equation

Yildirim OZDEMIR ¹, Mustafa ALP

¹ *Department of Mathematics, Duzce University, Duzce, Turkey*
E-mail: yildirimozdemir@duzce.edu.tr, alp.mustafa81@gmail.com

Abstract: The nonlocal boundary value problem for a parabolic-Schrödinger equation is considered. The stability estimates for the solution of the given problem are established. For approximately solving this nonlocal boundary problem the first and second order of difference schemes are presented. The theoretical statements for the solution of these difference schemes are supported by the result of numerical examples.

Keywords: finite difference equation, partial differential equation, stability

2010 Mathematics Subject Classification: 65N06, 65N12

REFERENCES

- [1] Ashyralyev, A., Fattorini, H. O., “On uniform difference scheme for second order singular perturbation problems in Banach spaces”, *SIAM J. Math. Anal.*, Vol.23, No.1, pp. 29-54, 1992.
- [2] Ashyralyev, A., Ozdemir, Y., “Stability of difference schemes for hyperbolic-parabolic equations”, *Comput. Math. Appl.*, Vol.50, pp. 1443-1476, 2005.
- [3] Ashyralyev, A., Ozdemir, Y., “On nonlocal boundary value problems for hyperbolic-parabolic equations”, *Taiwan. J. Math.*, Vol.11, No.4 pp. 1075-1089, 2007.
- [4] Ozdemir, Y., Kucukunal, M., “On nonlocal boundary value problems for hyperbolic Schrödinger equations”, *Abstr. Appl. Anal.*, Vol.2012, pp. 1-12, 2012.