

More general common fixed point theorems under a new concept

Hakima BOUHADJERA ¹

¹ *Laboratory of Applied Mathematics, Badji Mokhtar-Annaba University, Annaba, Algeria*

E-mail: b_hakima2000@yahoo.fr

Abstract: A general common fixed point theorem for two pairs of weakly subsequentially continuous mappings (recently introduced) satisfying a significant estimated implicit function is proved. An extension of this result is thereby obtained. Our results assert the existence and uniqueness of common fixed points in several cases.

Keywords: subcompatible mappings, R -weakly commuting mappings of type (\mathcal{A}_f) (resp. (\mathcal{A}_g)), sequentially continuous mappings of type (\mathcal{A}_f) (resp. (\mathcal{A}_g)), weakly reciprocally continuous mappings, weakly subsequentially continuous mappings.

2010 Mathematics Subject Classification: 47H10, 54H25, 47H06, 47H09.

REFERENCES

- [1] R.P. Agarwal, R.K. Bisht, N. Shahzad *A comparison of various noncommuting conditions in metric fixed point theory and their applications*, Fixed Point Theory and Applications (2014), 2014:38.
- [2] M.A. Al-Thagafi, N. Shahzad *Generalized I-nonexpansive selfmaps and invariant approximations*, Acta Math. Sin. 23 (2007), no. 6, 965-972.
- [3] H. Bouhadjera, C. Godet-Thobie *Common fixed point theorems for pairs of subcompatible maps*, arXiv: 0906.3159v1 [math. FA] 17 June 2009.
- [4] H. Bouhadjera, C. Godet-Thobie *Common fixed point theorems for pairs of subcompatible maps*, arXiv: 0906.3159v2 [math. FA] 23 May 2011.