

# Monotone operators and the proximal point algorithm in $CAT(0)$ spaces

G. Zamani Eskandani<sup>1</sup>, M. Raeisi<sup>2</sup>

<sup>1</sup> *Faculty of Sciences, Department of Mathematics, University of Tabriz,  
Tabriz, Iran*

*zamani@tabrizu.ac.ir*

<sup>2</sup> *Faculty of Sciences, Department of Mathematics, University of Tabriz,  
Tabriz, Iran*

*m.raeisi@tabrizu.ac.ir*

**Abstract:** In this paper, by using products of finitely many resolvents of monotone operators, we propose an iterative algorithm for finding a common zero of a finite family of monotone operators and a common fixed point of a finite family of nonexpansive mappings in  $CAT(0)$  spaces. We derive the strong convergence of the proposed algorithm under appropriate conditions.

**Keywords:** Proximal point algorithm,  $CAT(0)$  spaces, monotone operator

**2010 Mathematics Subject Classification:** 47H09, 47H10

## REFERENCES

- [1] B. A. Kakavandi, Weak topologies in complete  $CAT(0)$  metric spaces. Proc Am Math Soc. **141** ( 2013), 1029- 1039.
- [2] M. Bačák, Convex Analysis and Optimization in Hadamard Spaces, volume 22 of De Gruyter Series in Nonlinear Analysis and Applications. De Gruyter, Berlin, (2014).
- [3] S. Kamimura and W. Takahashi, Approximating solutions of maximal monotone operators in Hilbert spaces, J. Approx. Theory **13** (2000), 226—240.
- [4] R. T. Rockafellar, Monotone operators and the proximal point algorithm, SIAM J. Control Optim. **14** (1976), 877—898.