

A Second Order Difference Scheme for the Numerical Processing of Source Identification Terms in Hyperbolic Equations

Allaberen Ashyralyev¹, Fathi Emharab²

¹*Department of Mathematics, Near East University, Nicosia, TRNC, Mersin 10, Turkey.*

Peoples' Friendship University of Russia (RUDN University), Moscow 117198, Russia.

Institute of Mathematics and Mathematical Modeling, 050010, Almaty

allaberen.ashyralyev@neu.edu.tr

URL:<http://staff.neu.edu.tr>

²*Department of Mathematics, Near East University, Nicosia, TRNC, Mersin 10, Turkey.*

Department of Mathematics, Omar Al-Mukhtar University, El-Beida.

fathi.emharb@gmail.com

Abstract: In the present study, a second order difference scheme for a one-dimensional hyperbolic equation is presented. Stability estimates for the solution of the difference scheme are established. This difference scheme is tested on an example and some numerical results are presented.

Keywords: Source identification, hyperbolic differential equations, difference schemes, stability.

2010 Mathematics Subject Classification: 35J05, 35J08, 35J25

REFERENCES

- [1] A. Ashyralyev and A. U. Sazaklioglu, Investigation of a time-dependent source identification inverse problem with integral overdetermination, *Numerical Functional Analysis and Optimization* **38** (10):1276-1294, 2017.
- [2] M. Ashyralyeva and M. Ashyralyev, On a second order of accuracy stable difference scheme for the solution of a source identification problem for hyperbolic-parabolic equations, *AIP Conference Proceedings* 1759 (2016), 020023.
- [3] G. Di. Blasio, A. Lorenzi, Identification problems for parabolic delay differential equations with measurement on the boundary, *Journal of Inverse and Ill-Posed Problems*, **15**(7):709-734, 2007.