Numerical Solutions of Source Identification Problem for Parabolic-Elliptic Equations

A. Ashyralyev[1,2,3], Ç. Arıkan[1]

[1] *Department of Mathematics, Near East University, Nicosia, TRNC, Mersin 10, Turkey.*
[2] *Peoples' Friendship University of Russia (RUDN University), Moscow 117198, Russia.*
[3] *Institute of Mathematics and Mathematical Modeling, 050010, Almaty*

allaberen.ashyralyev@neu.edu.tr

URL:http://staff.neu.edu.tr
caginarikan@outlook.com

Abstract: A numerical study is presented of boundary value problem for parabolic-elliptic equations with unknown source term. The first and second order of accuracy difference schemes are constructed and the numerical procedure for implementation of these schemes is discussed. Numerical example is included.

REFERENCES

[1] A. K. Urinov, S. T. Nishonova, A problem with integral conditions for an elliptic-parabolic equation, *Mathematical Notes* 102 (1-2) (2017) 68-80.DOI: 10.1134/S0001434617070082.

[2] M. Kohlmann, On an elliptic-parabolic MEMS model with two free boundaries, *Applicable Analysis* 94 (10) (2015) 2175-2199. DOI: 10.1080/00036811.2015.1033410.

[3] R. Denk, T. Seger, L-p-estimates for a transmission problem of mixed elliptic-parabolic type., *Boundary Value Problems.* Article Number: 22 (2014). DOI: 10.1186/1687-2770-2014-22.

[4] I. C. Kim, N. Pozar, Nonlinear Elliptic-Parabolic Problems, *Arcive for Rational Mechanics and Analysis* 210(3) (2013) 975-1020. DOI: 10.1007/s00205-013-0663-3.

[5] A. Ashyralyev, On the well-posedness of the nonlocal boundary value problem for elliptic-parabolic equations, *Electronic Journal of Qualitative of Differential Equations* 49 (2011) 1-16.

[6] A. Ashyralyev, O.Gercek, Finite difference method for multipoint nonlocal elliptic-parabolic problems, *Computers and Mathematics with Applications* 60 (7) (2010) 2043-2052. DOI: 10.1016/j.camwa.2010.07.044.