

An application for time-fractional $K(m,m)$ model equations by symmetry method

Dogan Kaya¹, Gulistan Iskandarova²

Istanbul Commerce University, Faculty of Art and Science, Istanbul, Turkey

¹ *dkaya36@yahoo.com*

² *gulistan.iskandarova@gmail.com*

Abstract: In this work we consider the $K(m,m)$ model equations with time-fractional derivative [1]. We apply Lie symmetry group method for this equations, find infinitesimal operators and some exact solutions [3 - 5]. Also, we investigate conservation laws of this equations.

Keywords: Lie symmetry group, fractional derivative, conservation laws.

2010 Mathematics Subject Classification: 76M60, 26A33, 35L65

REFERENCES

- [1] P. Rosenau, J. Hyman, Compactons: solitons with finite wavelength, *Phys. Rev. Lett.* vol. 70, no. 5, 564–567, 1993.
- [2] L.V. Ovsiannikov, *Group analysis of differential equations*, Acad. Press, USA, 1982.
- [3] N. Ibragimov, *Lie group analysis classical heritage*, ALGA Publications Blekinge Institute of Technology Karlskrona, Sweden, 2004.
- [4] R.K. Gazizov, A.A. Kasatkin, S.Yu. Lukashchuk, Continuous transformation group of fractional differential equations. *Vestn. USATU.* vol. 9, 125–135, 2007.
- [5] G. Iskandarova and D. Kaya, Symmetry solution on fractional equation, *IJOCTA* vol. 7, no. 3, 255–259, 2017.