Distribution of eigenvalues of a third-order differential operator with strongly regular boundary conditions

Nurlan S. Imanbaev¹

¹ South-Kazakhstan State Pedagogical University, 13 Baytursinov str., 160012 Shymkent, Kazakhstan imanbaevnur@mail.ru

Abstract: This paper is devoted to the differential operator, generated by the ordinary differential equation and nonlocal boundary conditions. It is proved that the characteristic determinant of the considered spectral problem is an entire function.

The paper was supported by the state grant "The Best Teacher of the University - 2017" of the Ministry of Education and Science of the Republic of Kazakhstan

References

- G. D. Birkhoff, On the asymptotic chapacter of the solutions of certain linear differential equations containing a papameter, Transactions of the American Mathematical Society, Vol. 9, 2, 219–231 (1908).
- [2] G. D. Birkhoff, Boundary value and expansion problems of ordinary linear differential equations, Transactios of the American Mathematical Society, Vol. 9, 4, 373–395 (1908).
 [2] M. A. N. in the Linear Differential Quantum Mathematical Society, Vol. 9, 4, 1020 (1908).
- [3] M. A. Naimark, Linear Differential Operators, Moscow: Science, 496 p. (1969).
- [4] M. A. Sadybekov and N. S. Imanbaev, Reqular Differential Operator with Perturbed Boundary Condition, Mathematical Notes, Vol. 101, 5, 878–887. (2017).
- [5] N. S. Imanbaev, Stability of the basis property of eigenvalue systems of Sturm Liouville operators with integral boundary condition, Electron. J. Diff. Equ, Vol. 2016, 87, 1–8 (2016).